

Electronic supplementary materials

A new strategy for the design of macrocycles containing tetrahydrofuran pharmacophores with antitumor activity

V. A. D'yakonov,^{a} L. U. Dzhemileva,^{a*} I. I. Islamov,^b E. Kh. Makarova,^b
I. V. Gaisin,^b A. A. Makarov,^b U. M. Dzhemilev^a*

^a*N. D. Zelinsky Institute of Organic Chemistry Russian Academy of Sciences,
Leninsky prosp., 47, Moscow, 119991, Russia.*

E-mail: DyakonovVA@ioc.ac.ru

^b*Institute of Petrochemistry and Catalysis, Ufa Federal Research Center,
Russian Academy of Sciences, prosp. Oktyabrya, 141, Ufa, 450075, Russia*

NMR data




Figure S1. ^1H and ^{13}C NMR spectra of compound **3** in CDCl_3 .





Figure S2. ^1H and ^{13}C NMR spectra of compound 4 in CDCl_3 .





Figure S3. ^1H and ^{13}C NMR spectra of compound **5** in CDCl_3 .





Figure S4. ¹H and ¹³C NMR spectra of compound 6 in CDCl₃.





Figure S5. ¹H and ¹³C NMR spectra of compound **7** in CDCl₃.